

GALACTIC EMISSION AT 19 GHZ

ANGÉLICA DE OLIVEIRA-COSTA^{1,2,3}, MAX TEGMARK^{3,4}, LYMAN A. PAGE¹ & STEPHEN P. BOUGHN⁵

Accepted for publication in ApJL

ABSTRACT

We cross-correlate a 19 GHz full sky Cosmic Microwave Background (CMB) survey with other maps to quantify the foreground contribution. Correlations are detected with the Diffuse Infrared Background Experiment (DIRBE) 240, 140 and 100 μ m maps at high latitudes ($|b| > 30^\circ$), and marginal correlations are detected with the Haslam 408 MHz and the Reich & Reich 1420 MHz synchrotron maps. The former agree well with extrapolations from higher frequencies probed by the COBE DMR and Saskatoon experiments and are consistent with both free-free and rotating dust grain emission.

Subject headings: cosmic microwave background – methods: data analysis

1. INTRODUCTION

One of the major challenges in any Cosmic Microwave Background (CMB) anisotropy analysis is to determine the fraction of the observed signal due to diffuse Galactic emission. Three components of Galactic emission have been firmly identified: synchrotron and free-free radiation, which are important mainly at frequencies below 60 GHz, and thermal emission from dust particles, which is important mainly at frequencies above 60 GHz (see, *e.g.*, Weiss 1980; Bennett *et al.* 1992; Brandt *et al.* 1994; Tegmark and Efstathiou 1996). In principle, these three components can be discriminated by their frequency dependence and morphology. In practice, however, there is no component for which both the frequency dependence and the spatial template are currently well known (see, *e.g.*, Kogut *et al.* 1996a, hereafter K96a, and references therein).

The cross-correlation technique allows one to estimate the rms level of Galactic emission present in a CMB map. For instance, an analysis of high latitude Galactic emission in the COBE DMR map gave rms estimates of $(3.4 \pm 3.7) \mu\text{K}$ for synchrotron and $(2.7 \pm 1.3) \mu\text{K}$ for dust emission at 53 GHz on a 7° scale (Kogut *et al.* 1996b, hereafter K96b). A third component, correlated with the DIRBE maps but decreasing with frequency, was detected at the level of $(7.1 \pm 1.7) \mu\text{K}$ and tentatively identified as free-free emission (K96b). Although this component was also detected in the Saskatoon maps (de Oliveira-Costa *et al.* 1997, hereafter DOC97) on a 1° scale, at a level $(17.5 \pm 9.5) \mu\text{K}$ at 40 GHz, it is still not clear if it is due to free-free emission. An analysis of Owens Valley Radio Observatory (OVRO) data at 14.5 GHz showed such a component at the level of 203 μK in a small sky region on scales of $7' - 22'$ (Leitch *et al.* 1997, hereafter L97), but this level is substantially higher than free-free emission estimates based on H α images (Gaustad *et al.* 1996; Simonetti *et al.* 1996). Suggested explanations include the presence of high temperature gas (L97) and rotating dust grains (Draine & Lazarian 1998, hereafter DL98), but more data




FIG. 1.— 19 GHz survey (top), 408 MHz Haslam synchrotron template (middle), and DIRBE 100 μ m dust template (bottom).

is needed to settle this issue. The purpose of this letter is to evaluate the Galactic contribution to the 19 GHz full sky map (see Figure 1) by cross-correlating it with the DIRBE dust maps, and with the Haslam and Reich & Reich synchrotron maps.

2. METHOD

The 19 GHz map consists of $N = 24576$ pixels with sky temperatures y_i and noise n_i . We assume that this map is a superposition of CMB fluctuations and a Galac-

¹Princeton University, Dept. of Physics, Princeton, NJ 08544

²angelica@ias.edu

³Institute for Advanced Study, Olden Lane, Princeton, NJ 08540

⁴Hubble Fellow

⁵Haverford College, Dept. of Astronomy, Haverford, PA 19041

tic component whose angular distribution is traced in part by an external data set. Representing these contributions by N -dimensional vectors,

$$\mathbf{y} = \mathbf{n} + \mathbf{x}_{CMB} + \alpha \mathbf{x}_{Gal} + \mathbf{y}_{Gal}, \quad (1)$$

where x_{CMB}^i is the contribution of the fluctuating component of the CMB, x_{Gal}^i is the brightness fluctuations of the Galactic template map (not necessarily in temperature units), α is the coefficient that converts units of the Galactic template into antenna temperature and y_{Gal}^i represents any residual Galactic contribution which is uncorrelated with x_{Gal}^i . We consider $\mathbf{n}$ and $\mathbf{x}_{CMB}$ to be random variables with zero mean, i.e. $\langle \mathbf{x}_{CMB} \rangle = \langle \mathbf{n} \rangle = 0$, and $\mathbf{x}_{Gal}$ and $\mathbf{y}_{Gal}$ to be constant vectors. Thus the data covariance matrix is given by

$$\mathbf{C} \equiv \langle \mathbf{y} \mathbf{y}^T \rangle - \langle \mathbf{y} \rangle \langle \mathbf{y}^T \rangle = \langle \mathbf{x}_{CMB} \mathbf{x}_{CMB}^T \rangle + \langle \mathbf{n} \mathbf{n}^T \rangle, \quad (2)$$

where $\langle \mathbf{x}_{CMB} \mathbf{x}_{CMB}^T \rangle$ is the covariance matrix of the CMB and $\langle \mathbf{n} \mathbf{n}^T \rangle$ is the noise covariance matrix. The noise in the 19 GHz map is approximately uncorrelated and has an rms amplitude of $\sigma_i \sim 2$ mK. Therefore, the covariance matrix of this map is

$$C_{ij} \approx \langle n_i n_j \rangle \approx \sigma_i^2 \delta_{ij}. \quad (3)$$

Minimizing $\chi^2 \equiv (\mathbf{y} - \alpha \mathbf{x}_{Gal})^T \mathbf{C}^{-1} (\mathbf{y} - \alpha \mathbf{x}_{Gal})$ yields the minimum-variance estimate of α , i.e.

$$\hat{\alpha} = \frac{\mathbf{x}_{Gal}^T \mathbf{C}^{-1} \mathbf{y}}{\mathbf{x}_{Gal}^T \mathbf{C}^{-1} \mathbf{x}_{Gal}} \quad (4)$$

with variance

$$\Delta \hat{\alpha}^2 = \frac{1}{(\mathbf{x}_{Gal}^T \mathbf{C}^{-1} \mathbf{x}_{Gal})}. \quad (5)$$

Note that unlike the case in dOC97, there is little contribution from chance alignments between the CMB and the various template maps, since the CMB contribution to $\mathbf{C}$ is negligible. If the noise is correlated or the anisotropy signal is significant, then our $\mathbf{C}$ used above will differ from the true covariance matrix, denoted $\mathbf{C}'$. Although eq. (4) still provides a reasonable and unbiased estimate of α^1 , its variance will be larger than implied by eq. (5), given by

$$\Delta \hat{\alpha}^2 = \frac{(\mathbf{x}_{Gal}^T \mathbf{C}^{-1} \mathbf{C}' \mathbf{C}^{-1} \mathbf{x}_{Gal})}{(\mathbf{x}_{Gal}^T \mathbf{C}^{-1} \mathbf{x}_{Gal})^2}. \quad (6)$$

In the next section, an estimate of $\mathbf{C}'$ is made from the data and the corresponding variance of $\hat{\alpha}$ is evaluated.

3. DATA ANALYSIS AND RESULTS

The 19 GHz map has an angular resolution of 3° FWHM and is stored in $1.3^\circ \times 1.3^\circ$ pixels (Cottingham 1987; Boughn *et al.* 1992). The template maps are convolved with a 3° Gaussian beam and regions within 20° and 30° of the Galactic plane are excluded. To avoid contamination by zodiacal dust emission, data within 10° of the Ecliptic plane are also excluded from the analysis; although, the results are found to be independent of this cut. Off

the Galactic plane, the 19 GHz map is dominated by the CMB dipole ($\ell=1$). On the other hand, because of its planar structure, emission associated with the Galaxy has a strong quadrupole ($\ell=2$) component. Therefore, the monopole, dipole, and quadrupole moments are removed from both the 19 GHz and template maps. As a consequence, the computed $\hat{\alpha}$ depends only on correlated structure in the maps on angular scales $\lesssim 90^\circ$.

3.1. Correlations and their Error Bars

The 19 GHz map was cross-correlated with five different templates: two for synchrotron emission, the 408 MHz (Haslam *et al.* 1981) and 1420 MHz (Reich and Reich 1988) surveys; and three to study dust and free-free emission, the 100, 140 and $240\mu\text{m}$ Diffuse Infrared Background Experiment (DIRBE) sky maps (Boggess *et al.* 1992). Table 1 lists the coefficients $\hat{\alpha}$ derived from eq. (4) with errors computed from eq. (5). All three DIRBE templates show significant correlations with the 19 GHz map, while the two synchrotron templates are found to be only marginally correlated.

Also listed in Table 1 are the implied fluctuations in antenna temperature in the 19 GHz map, i.e. $\delta T = \hat{\alpha} \sigma_{Gal}$, where σ_{Gal} is the rms of the template map. If a template map includes a distinct component that is uncorrelated with the 19 GHz map, then δT is underestimated by a factor $\sigma'_{Gal}/\sigma_{Gal}$, where σ'_{Gal} is the rms of the correlated component of the template map. For this reason, the δT 's in Table 1 should be considered lower limits.

As mentioned above, the errors listed in Table 1 were computed from eq. (5) and are therefore lower limits to the error. We now describe a series of tests, performed to estimate the uncertainty in $\hat{\alpha}$ due to correlated noise and other systematics.

3.1.1. The Noise Correlation Function

If the true noise correlation matrix is isotropic, $C'_{ij} = \langle n_i n_j \rangle = R(\theta_{ij}) \sigma_i \sigma_j$ where θ_{ij} is the angle between pixels i and j , then we can estimate the noise correlation function $R(\theta)$ of the 19 GHz data (after removing the monopole, dipole, and quadrupole) by $R(\theta) = N_\theta^{-1} \sum_{ij} y_i y_j / \sigma_i \sigma_j$, where y_i is the antenna temperature of the i^{th} pixel and the sum is over all N_θ pairs of pixels separated by θ . Substituting these relations into eq. (6) gives estimates of $\Delta \hat{\alpha}$ which are from 6% to 17% larger than those in Table 1.

3.1.2. Monte Carlo Simulations

As another test of the robustness of the estimates of $\hat{\alpha}$, we employed Monte Carlo simulations in which the template maps are sliced into eighteen regions of equal area, each corresponding to a range of Galactic latitude $|b|$. Inside each of these regions the pixels are rearranged in random order, so that the latitude dependence is preserved but the longitudinal correlations are destroyed. Repeating this procedure 1000 times yields distributions of $\hat{\alpha}$'s consistent with zero mean and with standard deviations $\Delta \hat{\alpha}_M \sim 1.6$ times larger than the formal errors $\Delta \hat{\alpha}$ (see Table 1).

¹Note that the diagonal approximation of eq. (3) reduces eq. (4) to a simple noise-weighted least-squares fit.

3.1.3. Sky Rotations

Because of the approximate axial symmetry of the Galaxy, it is natural to ask if the correlations are simply due to overall large-scale Galactic structure common to all emission components. To test this hypothesis, we repeated the analysis with $2 \times 2 \times 36 = 144$ transformed maps, rotated around the Galactic axis by multiples of 10° and/or flipped vertically and/or horizontally.

For a 20° Galactic cut, the correct template map has the highest of all 144 correlations, while the distribution of these correlations has standard deviation $\Delta\hat{\alpha}_T \sim 2.5$ times larger than the formal errors $\Delta\hat{\alpha}$. Likewise, the correct DIRBE maps have the highest of all 144 correlations and the standard deviations are $\Delta\hat{\alpha}_T \sim \Delta\hat{\alpha}_M$, even for a 30° Galactic cut (see Table 1). In contrast, we find no significant correlation between the $100\mu\text{m}$ and Haslam maps, indicating that synchrotron and dust emission are not strongly correlated at high latitudes.

Since both $\Delta\hat{\alpha}_M$ and $\Delta\hat{\alpha}_T$ have correlated signal contributing to the noise, they constitute overestimates of the true error bars.

3.2. Latitude Dependence

To investigate the dependence of the correlation on Galactic latitude, we sliced the maps into six regions of equal area, each corresponding to a range of latitude $|b|$. Figure 2 shows the results for the $100\mu\text{m}$ map. Note that $\hat{\alpha}$ from the 19 GHz DIRBE correlation is almost the same at each latitude band, indicating that this correlation is not dominated by one or two nearby clouds. The 19 GHz Haslam correlation is found to be more concentrated in the Galactic plane².




FIG. 2.— The dependence of $\hat{\alpha}$ on Galactic latitude $|b|$ for the $100\mu\text{m}$ map. The error bars are given by $\Delta\hat{\alpha}_T$.

3.3. Scale Dependence

One way to determine angular scale of the correlation is to compute the angular cross-correlation function (CCF) of the 19 GHz map with the template maps. We computed this according to $CCF(\theta) = [\sum_{ij} x_{Gal,j} y_i / \sigma_i^2] / [\sum_{ij} \sigma_i^{-2}]$, where the sums are again over all pairs of pixels separated by θ .

As an example, Figure 3 shows the $CCF(\theta)$ of 19 GHz and $140\mu\text{m}$ map, as well as the auto-correlation function of $140\mu\text{m}$ map. It is clear that the correlated structure in

the two maps is on small angular scales ($\theta \lesssim 10^\circ$) and that the CCF is well-behaved on all angular scales. The latter gives us additional confirmation that there are no large, unknown systematics that compromise the analysis. Similar results are found for the other two DIRBE maps. The lower signal-to-noise for the correlation of the two radio templates result in CCF's which are less demonstrative. Note that $CCF(0) \propto \hat{\alpha}$.




FIG. 3.— $CCF(\theta)$ between 19 GHz and $140\mu\text{m}$ map for a 30° Galactic cut, together with the $140\mu\text{m}$ auto-correlation.

As another way to investigate the dependence of the correlation on the angular scale, we repeated the analysis after high-pass filtering the 19 GHz map using a partial sky multipole removal technique (see Tegmark & Bunn 1995). Only after removing spherical harmonic components with $\ell \gtrsim 8$ does the correlation significantly decrease, which again indicates that large-scale Galactic structure is not the source of the correlation. Both types of filtering therefore indicate that the bulk of the correlations are caused by fluctuations on scales of several degrees, which is consistent with the shape of the CCF in Figure 3.

4. CONCLUSIONS

The two synchrotron templates are found to be marginally correlated with the 19 GHz map, while all three DIRBE far-infrared templates show a significant correlation. When a synchrotron and a DIRBE template are simultaneously fit to the 19 GHz map (via linear regression) the correlation coefficients do not change significantly from those listed in Table 1 and, in addition, the two fit parameters are essentially uncorrelated ($|\rho| \lesssim 0.05$). We conclude that the correlations with DIRBE dust emission are independent of the correlations with synchrotron emission. The amplitude of the signal is much larger than expected for ordinary (vibrational) dust emission, as shown in Figure 4. Moreover, there is now good agreement between different experiments that this correlated component is brighter at lower frequencies.

So what physical component is this? Two contenders have been proposed. K96ab argue on physical grounds that free-free emission might be spatially correlated with dust. However, the correlations between $\text{H}\alpha$ (which is normally a good tracer of free-free emission) and CMB maps, and between $\text{H}\alpha$ and the DIRBE maps, are weak (L97;

²In order to test how important spatially localized features are for the correlation, we cross-correlated 19 GHz with Haslam using only one hemisphere. Most of the correlation was found to come from the northern hemisphere, presumably because of emission features such as the North Polar Spur.




FIG. 4.— Frequency dependence of the fluctuating component of the Galactic emission for $|b| > 30^\circ$. The DMR correlations are represented by filled circles (K96b), Saskatoon data by pentagons (dOC97) and 19 GHz by a square with error bar given by $\Delta\hat{\alpha}_T$ (this work). Dust emission (vibrational modes only) is normalized using the DMR correlation at 90 GHz (K96a). The free-free model is normalized to the DMR correlation at 31.5 GHz (the rms free-free emission derived from H α is much smaller than this DIRBE-correlated component). The synchrotron emission is normalized to the correlation between the 19 GHz map and the 408 MHz Haslam map (see Table 1), and is consistent with the 31.5 GHz upper limits from K96b. The thickness of each curve corresponds to the normalization uncertainty. The dashed line is an example of emission from rotating dust grains (DL98). All fluctuations have been converted to DMR angular scales as $\delta T = \varepsilon \hat{\alpha} \sigma_{Gal}^i$, where σ_{Gal}^i is the rms of each template map and the correction factor is

$$\varepsilon = \left(\left[\sum (2\ell + 1) \ell^\beta W_\ell^{DMR} \right] / \left[\sum (2\ell + 1) \ell^\beta W_\ell^i \right] \right)^{1/2}.$$

Here W_ℓ^{DMR} is DMR window function, W_ℓ^i is the window function of the experiment to be converted and we assume a $\beta = -3$ power spectrum slope.

Kogut 1997; McCullough 1997). One possibility is the presence of an extremely hot ($\gtrsim 10^6$ K), ionized plasma (L97); however, DL98 have argued that this cannot be true across the whole sky on energetic grounds. These results motivated DL98 to suggest that perhaps it is dust after all, but emitting through rotational rather than vibrational excitations. As shown in Figure 4, this can give

a spectrum quite similar to that of free-free emission in the relevant frequency range.

H α maps with better accuracy are currently being made by the Dennison, Gaustad and Reynolds groups and should help to settle the issue. Another test is indicated by Figure 4: although both the rotating dust and the free-free models appear consistent with the available data³, the former predicts a bump and a downturn around 10–20 GHz while the latter predicts a continued rise toward lower frequencies. A cross-correlation analysis with lower frequency data like the 10 GHz Tenerife map may be able to discriminate between these two models.

We would like to thank Ed Cheng, Dave Cottingham, Bruce Draine, Dale Fixsen, Ken Ganga, Ed Groth, Al Kogut, Alex Lazarian, George Smoot and David Wilkinson for helpful comments. Support for this work was provided by NASA grants NAG5-6034 and NAG5-3015, by NSF grant PHY-9600015, by a David & Lucile Packard Foundation Fellowship (to LP), and by NASA through Hubble Fellowship HF-01084.01-96A from STScI, operated by AURA, Inc. under NASA contract NAS5-26555.

Table 1 – Correlations with the 19 GHz map

Template ^(a)	$\hat{\alpha}^{(b)}$	$\Delta\hat{\alpha}$	$\Delta\hat{\alpha}_M$	$\Delta\hat{\alpha}_T$	$\delta T^{(c)}$ (μ K)
100 μ m	38.5	3.3	5.8	8.3	138.6 $\pm$ 29.9
140 μ m	29.8	2.5	4.0	6.3	146.0 $\pm$ 30.9
240 μ m	46.2	3.9	6.0	10.0	143.2 $\pm$ 31.0
408 MHz	13.4	3.1	3.4	4.4	52.3 $\pm$ 17.2
1420 MHz	0.9	0.1	0.2	0.3	86.5 $\pm$ 28.8
100 μ m	47.1	9.0	16.2	17.3	65.9 $\pm$ 24.2
140 μ m	31.6	6.3	10.2	10.3	66.4 $\pm$ 21.6
240 μ m	38.9	8.4	12.8	12.9	66.1 $\pm$ 21.9
408 MHz	7.6	3.7	4.7	6.2	25.1 $\pm$ 20.5
1420 MHz	0.4	0.2	0.2	0.3	29.8 $\pm$ 22.3

^(a) Correlations for $|b| > 20^\circ$ (top) and $|b| > 30^\circ$ (bottom).

^(b) $\hat{\alpha}$ has units μ K (MJy/sr)⁻¹ for the DIRBE templates, μ K/K for the 408 MHz template and μ K/mK for the 1420 MHz template.

^(c) $\delta T \equiv (\hat{\alpha} \pm \Delta\hat{\alpha}_T) \sigma_{Gal}$.

REFERENCES

- Bennett, C.L., *et al.* 1992, ApJ, 396, L7
 Boggess, N.W., *et al.* 1992, ApJ, 397, 420
 Boughn, S.P. *et al.* 1992, ApJ, 391, L49
 Brandt, W.N., *et al.* 1994, ApJ, 424, 1
 Cottingham, D.A. 1987, Ph.D. thesis, Princeton University.
 de Oliveira-Costa, *et al.* 1997, ApJ, 482, L17 (dOC97)
 Draine, B.T., Lazarian, A. 1998, ApJ, 494, L19 (DL98)
 Gaustad, J.E., *et al.* 1996, PASP, 108, 351
 Haslam, C.G.T., *et al.* 1981, A&A, 100, 209
 Kogut, A., *et al.* 1996a, ApJ, 460, 1 (K96a)
 Kogut, A., *et al.* 1996b, ApJ, 464, L5 (K96b)
 Kogut, A. 1997, AJ, 114, 1127
 Leitch, E.M., *et al.* 1997, ApJ, 486, L23 (L97)
 McCullough, P.R. 1997, AJ, 113, 2186
 Reich, P., Reich, W. 1988, A&AS, 74, 7
 Simonetti, J.H., *et al.* 1996, ApJ, 458, L1
 Tegmark, M., Bunn, E. 1995, ApJ, 455, 1
 Tegmark, M., Efstathiou, G. 1996, MNRAS, 281, 1297
 Weiss, R. 1980, ARAA, 18, 489

³We compute the spectral index between 19 and DMR 53 GHz to be ~ -2.4 , and between 19 GHz and 1420 MHz to be ~ -2.8 .